

Ecosystem-based Adaptation for Increased Water Availability: A Case Study from the Hills and Mountains of Nepal

Keshav Prasad Khanal^{1*}, Top Bahadur Khatri¹, Santosh Mani Nepal¹,
Buddi Sagar Poudel², Raju Sapkota², Binod Thapa¹,
Digambar Singh Dahal¹ and Bhola Dhakal¹

¹*Ecosystem-based Adaptation Project II, Kathmandu, Nepal,*

²*Ministry of Forests and Environment, Kathmandu, Nepal*

*Corresponding Author: Keshav_khanal@hotmail.com

Abstract

Water scarcity is a pressing issue for Nepal's rural hill communities, worsened by climate change, irregular rainfall and unplanned infrastructure. This study examines the impact of Ecosystem-based Adaptation (EbA) strategies on water availability and soil moisture in Dolakha, Salyan and Achham districts. Using a participatory approach, data were collected through household surveys and focus group discussions to assess community perceptions of EbA interventions, particularly construction and rehabilitation of conservation ponds. Results show that EbA measures significantly improve water availability and soil moisture, with perception scores ranging from 6.12 to 6.84 on a 10-point scale. Additionally, reduced time spent on collecting water benefits women and children, improving their quality of life. These findings underscore the importance of integrating EbA strategies into government policies and development programmes, providing sustainable solutions to water scarcity and climate variability while enhancing community resilience.

Keywords: Climate change, ecosystem-based adaptation, water conservation

INTRODUCTION

Nepal is one of the most climate change-vulnerable countries in the world. In response to this, the country has been reforming policies and implementing climate change mitigation and adaptation strategies. Key initiatives include the National Climate Change Policy 2019, the National Adaptation Plan for 2021–2050 (MoFE 2023), the Second Nationally Determined Contributions (NDC) and gender-responsive climate-resilient adaptation plans (MoFE 2020).

Irregular rainfall patterns in Nepal are severely impacting the livelihoods of farmers, particularly in rural areas. Water

shortages are prevalent during the pre-monsoon season (March–May) and the post-monsoon and winter seasons (October–February) (GFDRR 2021). To address vulnerability to these erratic rainfall patterns, there is a pressing need for establishing a year-round water supply system. Water conservation ponds are an effective adaptation strategy as they capture rainwater, replenish groundwater reserves during the monsoon, and help reduce soil erosion and surface runoff, especially on landslide-prone slopes (Koirala 2021). Moreover, these ponds ensure water availability during dry periods, which is vital for traditional rural livelihoods, like

livestock farming and crop cultivation. They also support new activities, such as fish farming, and help stabilise and revegetate gullies, enabling small-scale cultivation of fodder, vegetables and fruit trees along the pond banks. Additionally, conservation ponds reduce the time women spend on fetching water (FAO 2014).

Springs and wells (*Kuwa*) are crucial water sources in the hills and mountains of Nepal. In the Hindu Kush Himalaya (HKH) region, temperatures are projected to rise by more than 2°C on average by 2050 (Shrestha *et al.* 2019). Nepal is predominantly mountainous, with about 43 per cent of its total land area classified as hills. The varied landscapes in these hilly regions create niche microclimatic habitats that support diverse ecological zones and farming systems (Shrestha *et al.* 2015).

Rural areas in the hills and mountains face significant challenges regarding water availability for drinking and irrigation. Factors such as unplanned infrastructure development, changes in land use practices, prolonged droughts and erratic rainfall due to climate change have contributed to decreased water yield. Traditional knowledge-based water management practices, which include the use of local resources and techniques, like water conservation ponds, bioengineering methods for water source protection and tree planting near water sources to enhance yield and control erosion, have declined over the past decades. There could be many reasons for the decline, but the major ones could be the lack of intergenerational knowledge transfer, outmigration from rural areas to urban centres (Koirala 2021), changes in community structures and decreasing collective efforts.

The socioecological systems refer to the interconnected and dynamic interactions

between human societies and their natural environment. These systems encompass the way communities in the hills interact with, depend on and manage natural resources while adapting to environmental, social and economic changes. To adapt to climate adversities and build resilience in this socioecological systems, local communities have initiated various ecosystem-based adaptation (EbA) approaches for water management (CGED–Nepal 2024). These include promoting efficient water use through harvesting, storage, source protection and efficient usage. Several EbA measures have proven effective in building the adaptive capacity of local communities by offering simple and affordable technologies.

The EbA II project was implemented from March 2019 to April 2025 to reduce climate vulnerability in Nepal. The project was led by the Government of Nepal, Ministry of Forests and Environment (MoFE), Global Environment Facility (GEF) and United Nations Environment Programme (UNEP) and funded by the Least Developed Countries Fund (LDCF). It implemented EbA measures to restore forests and rangelands, mainly in mid-hills (Achham and Salyan) and high hills (Dolakha), reducing community sensitivity to climate change. The project adopted the approach of integrating ecosystem-based solutions to address climate change impacts, while enhancing biodiversity conservation and improving community resilience. The project emphasised the role of natural ecosystems in mitigating the effects of climate change, such as floods, landslides and droughts. By restoring and protecting ecosystems like forests, wetlands and watersheds, EbA aimed to reduce vulnerability and enhance adaptive capacities of local communities, especially in marginalised and vulnerable regions. Water conservation was one of the main achievements of the project.



This paper provides evidence of the efficiency of the EbA measures, such as recharge ponds and conservation ponds, in terms of increased water availability and infiltration in the EbA II project, implemented in Dolakha, Salyan and Achham districts in Bagmati, Karnali and Sudur Paschim provinces respectively, using a participatory approach. Proper documentation of how these measures improve water yield is essential for replicating them elsewhere. This study aims to assess and analyse the effectiveness of the EbA measures in enhancing water availability and yield, as well as their socioeconomic benefits for rural communities in the hills and mountains of Nepal. The research provides evidence-based insights into the impact of these measures and to generate recommendations for scaling up and integrating EbA practices into the local and national climate adaptation strategies.

Objectives of the study

The general objective of this study was to assess the impact of the EbA interventions on water availability and soil moisture improvement in the EbA II project areas.

The specific objectives of the study were: (i) to evaluate the impact of the EbA interventions on water availability in the spring sheds in Dolakha, Salyan and Achham districts; (ii) to assess the improvement in soil moisture levels resulting from water conservation and management measures under the EbA interventions; and (iii) to analyse community perceptions of the changes in water availability and soil moisture in the targeted areas.

STUDY METHODOLOGY

Study area

For this study, five conservation ponds and their beneficiary communities from Dolakha and Salyan districts each and six from Achham district were randomly selected. The list of the total number of conservation ponds constructed in the EbA II project sites by years is provided in Table 1. The EbA II project supported local communities in constructing and rehabilitating 92 conservation and water recharge ponds across the project districts (Table 1).

Table 1: Water conservation ponds constructed by EbA II project

District	Unit	2021	2022	2023	Total	Sample
Dolakha	Number	6	12	6	24	5
Salyan	Number	7	7	5	19	5
Achham	Number	18	21	10	49	6
Total		31	40	21	92	16

The list of water conservation ponds and communities selected for detailed household survey is presented in Table 2.

Table 2: List of water conservation ponds and communities selected for study

District	Municipality	Ward No.	Place
Dolakha	Shailung	4	Mathillo Gaon Jogidanda
	Shilung	4	Pokharidanda
	Bhimeshwor	9	Mane
	Kalinchowk	6	Okhreni
	Kalinchowk-5	5	Bandethali
	Kumakh	2	Danda Pipal Sim Narayan Takura
Salyan	Kumakh	2	Kiya Khola, Danda Kateri
	Bangaad Kupinde	1	Samaila
	Bangaad Kupinde	5	Sano Barule
	Bangaad Kupinde	7	Deutipujne
	Sanfebagar	13	Babla
Achham	Ramaraoshan	5	Badapani
	Ramaraoshan	6	Basanta, Bhatakaatiya
	Ramaraoshan	6	Sherapatalkharka, Bhatakaatiya
	Mellekh	6	Jhadigau, Sodsha
	Mellekh	6	Dudhimela, Sodsha

Sample size

Out of the 92 water conservation ponds constructed with the support of the EbA II project, 16 ponds were randomly selected: five each in Dolakha and Salyan districts and six in Achham district. This selection was made to ensure geographic diversity and manage logistical and resource constraints while maintaining statistical relevance for the study.

Research methodology

Before initiating data collection, an extensive literature review was conducted.

This review primarily included national policies, programmes and archival research. The research methodology employed both quantitative (household surveys) and qualitative (FGD and key informant interview) methods.

The survey was conducted in October 2023. A total of 392 households were surveyed across the three districts using pre-designed questionnaires, representing 10 per cent of the total beneficiaries (3,920) of the EbA II project. Table 3 summarises the proportional distribution of surveyed households across the three districts, ensuring a representative sample.

**Table 3: Distribution of surveyed households**

District	Total beneficiaries	Surveyed households	Percentage surveyed
Dolakha	1250	125	10
Salyan	1260	126	10
Achham	1410	141	10
Total	3920	392	10

Additionally, 16 FGDs were conducted. The household survey aimed to gather local perceptions of the effectiveness of the EbA interventions in increasing the water yield and soil moisture. The questionnaire included a variety of multiple-choice and open-ended questions.

To avoid interviewer bias and to maintain a cost-effective approach, a self-administered questionnaire was used. For data analysis and reporting, both quantitative and qualitative data were triangulated to provide a comprehensive analysis. Microsoft Excel was used for data entry and analysis.

We applied basic statistical methods, such as calculating the mean, standard deviation and confidence interval (CI). The CI test was used to estimate the range within which the true population mean is likely to fall, with 95 per cent confidence. This was done using the formula provided by Triola (2020) for cases where the population standard deviation is unknown.

Equation 1

$$CI = \bar{x} \pm z \cdot \frac{s}{\sqrt{n}}$$

Where:

\bar{x} is the sample mean,

z is the critical value from the standard normal distribution (1.96 for a 95% confidence level),

s is the sample standard deviation,

n is the sample size.

RESULTS

Major water-related problems in the study sites

The household survey and FGDs revealed that climate change poses numerous challenges related to water resources in the study areas. The major problems identified were:

1. **Dry spells:** The precipitation patterns have changed in the study areas. The altered patterns have resulted in, among other effects, more frequent and severe dry spells. These shifts have led to critical issues such as drying water sources and water scarcity, which have, in turn, significantly impacted agriculture, drinking water supplies and ecosystems.
2. **Water scarcity:** There are noticeable changes in precipitation patterns, with prolonged droughts leading to significant water scarcity.
3. **Floods:** More intense rainfalls are increasing flooding, leading to frequent sediment deposition in the downstream of watershed.
4. **Changes in water availability:** Scarcity of irrigation water during the crop cultivation period has led to a decline in agricultural productivity.

Community contribution to water conservation and management

Community contributions are crucial for achieving sustainable water conservation and management goals, as they leverage local knowledge, resources and collective action to effectively address water-related challenges. The participants reported their involvement in various water conservation and management activities. Their key contributions include constructing conservation ponds, regularly maintaining existing water sources, building water storage tanks, planting vegetation near water sources, maintaining water channels and protecting water sources.

Benefits of water conservation and management activities

Water conservation activities in the study areas are benefiting various stakeholders. The participants noted that local communities were the primary beneficiaries of these efforts, as they helped ensure reliable access to clean water for drinking, sanitation and household uses, especially during dry seasons when water availability may be scarce. They also reported improved agricultural productivity and significant enhancement of livelihoods of many community members.

Furthermore, 71 per cent of respondents reported that protecting and conserving water sources has positively impacted ecosystems and wildlife. Similarly, 62 per cent of respondents mentioned that women and girls, who are responsible for collecting water for drinking, cooking, washing,

hygiene and livestock raising, have received direct benefits from these conservation activities, leading to a significant reduction in their workload.

Programme sustainability

When asked about the programme's sustainability, 71.5 per cent of respondents indicated that the community was willing to voluntarily construct additional water conservation ponds. They also planned to contribute to the maintenance of existing ponds, expansion of pond sizes and protection of other water sources. Additionally, other members of the community, such as livestock herders, will apply their acquired knowledge and participate in water conservation efforts. Local communities also expressed their intention to coordinate with rural or municipal authorities to incorporate additional ponds, catchment protection activities and budget allocations into their annual development programmes.

Perception of local people on the increase of water availability and soil moisture following the construction of conservation pond

Dolakha

Out of the total 125 respondents, 85 (68%) responded that the community was willing to voluntarily construct additional water conservation ponds.




Figure 1: Perception of local people on the increase of water availability following the construction of conservation ponds in Dolakha district

The household survey shows that the residents believe there has been a significant improvement in water availability (at 95% confidence level) following the construction of conservation ponds. When asked to rate the improvement in water yield on a scale of 1 to 10, with 1 indicating no improvement and 10 indicating very good improvement, the average score for 125 participants was 6.48, with a standard deviation of ± 0.062 . While calculating the CI using equation 1, the CI ranged from 6.469 to 6.491. Both limits

exceed the mid value of 5 on a 1 to 10 scale. Thus, the result demonstrates a statistically significant positive community perception of improvement in water availability.

Similarly, when asked about the rating of improvement in greenery on a scale of 1 to 10, with 1 indicating no improvement and 10 indicating very good improvement, the average score for the 125 participants was 6.405, with a standard deviation of ± 1.338 . While calculating the CI using equation 1, the




Figure 2: Perception of local people on the improvement of greenery following the construction of conservation ponds in Dolakha district

CI ranged from 6.170 to 6.640. Both limits exceed the mid value of 5 on a 1 to 10 scale. Thus, the result demonstrates a statistically significant positive perception of local people on improvement in soil moisture following the construction of conservation ponds.

Salyan

A total of 126 community members shared their perceptions of water availability in the village following the implementation of various activities. On a scale of 0 to 10, the

average score given by the respondents was 6.12, with a standard deviation of ± 1.413 . This indicates that the local communities perceived a significant increase in water yield in the watershed following the construction of conservation ponds. While calculating the CI using equation 1, the CI ranged from 5.873 to 6.367. Both limits exceed the mid value of 5 on a 1 to 10 scale. Thus, the result demonstrates a statistically significant positive perception of improvement in water yield after the construction of conservation ponds.




Figure 3: Perception of local people on the increase of water yield following the construction of conservation ponds in Salyan district

Similarly, when asked regarding the improvement in greenery on a scale of 1 to 10, with 1 indicating no improvement and 10 indicating very good improvement, the average score for the 126 participants was 6.14, with a standard deviation of ± 1.289 . While calculating the CI using equation

1, the CI ranged from 5.915 to 6.365. Both limits exceed the mid value of 5 on a 1 to 10 scale. Thus, the result demonstrates a statistically significant positive perception of improvement in soil moisture following the construction of conservation ponds.




Figure 4: Perception of local people on the improvement of greenery after the construction of conservation pond in Salyan district

Achham

A total of 141 community members shared their perceptions of water availability in the village after various activities were implemented. On a scale of 0 to 10, the average score given by respondents was 6.84, with a standard deviation of ± 1.441. While

calculating the CI using equation 1, the CI ranged from 6.602 to 7.078. Both limits exceeded the mid value of 5 on a 1 to 10 scale. Thus, the result demonstrates a statistically significant positive perception of improvement in water availability following the construction of conservation ponds.




Figure 5: Perception of local people on the increase of water yield following the construction of conservation pond in Achham district

Similarly, when asked to rate the improvement in soil moisture and greenery on a scale of 1 to 10, with 1 indicating no improvement and 10 indicating very good improvement, the average score from the 141 participants was 6.84, with a standard deviation of ± 1.418 . While calculating the CI using equation

1, the CI ranged from 6.610 to 7.075. Both limits exceed the mid value of 5 on a 1 to 10 scale. Thus, the result demonstrates a statistically significant positive perception of improvement in soil moisture following the construction of conservation ponds.



Figure 6: Naula Samrrakshan Pokhari of ward no. 13, Babla, Sanfebagar Municipality, Achham



Figure 7: Dudhimela Conservation Pond of Sodasa, Mellekh-6, Achham




Figure 8: Perception of local people on the improvement of greenery after the construction of conservation pond in Achham district



DISCUSSION

The findings of this study underscore the critical role of the EbA interventions in enhancing water availability and soil moisture in the hill and mountain districts of Nepal. This discussion interprets the results in the context of the general and specific objectives, providing insights into the impacts, challenges and future implications of the interventions.

The general objective of assessing the impact of the EbA interventions on water availability and soil moisture improvement was met through a combination of quantitative and qualitative data collection methods. The data indicate that the EbA measures, particularly the construction and rehabilitation of conservation ponds, have positively influenced water resource availability and soil moisture levels. This aligns with global evidence on the effectiveness of nature-based solutions in addressing water scarcity and climate variability (Garcia-Herrero 2022).

Improvement in water availability

The household survey and FGDs revealed that the construction of conservation ponds has significantly enhanced water availability in the springsheds of Dolakha, Salyan and Achham districts. The average perception scores for water availability improvements ranged from 6.12 in Salyan to 6.84 in Achham, reflecting widespread community acknowledgment of increased water yield. The findings are like those of Thiaw (2013), which demonstrated that EbA interventions helped mitigate the impacts of erratic rainfall and prolonged dry spells, thereby ensuring a reliable water supply for drinking, irrigation and other livelihood activities.

Improvement in soil moisture

The study also highlights the role of the EbA measures in improving the soil moisture level. The perception scores for soil moisture and greenery improvements were consistent across the districts, with Achham reporting the highest average score of 6.84. Enhanced soil moisture has resulted in increased agricultural productivity, improved vegetation cover and reduced erosion, all of which contribute to the sustainability of local ecosystems. These findings are similar to those of Liniger *et al.* (2011). These benefits underscore the importance of integrating soil and water conservation strategies into adaptation planning.

Community perceptions of EbA benefits

Community perceptions provide critical insights into the socioeconomic impacts of the EbA interventions. The participants reported reductions in the time and effort required to collect water, particularly benefiting women and children. This finding is in line with the report of the Secretariat of the Convention on Biological Diversity (SCBD 2009), which highlights the social benefits of the EbA interventions, including improved water availability and reduced burden on women and children responsible for water collection in many regions. FGDs emphasised the broader benefits of conservation ponds, including increased biodiversity, enhanced groundwater recharge and stabilisation of gullies. This finding is consistent with the findings of the report, *Building a common vision for sustainable food and agriculture: Principles and approaches* (FAO 2014). This report highlights the contribution of water conservation practices such as pond construction to ecosystem

services, like biodiversity enhancement, water resource management and land stability. The community's willingness to contribute to the maintenance and expansion of these interventions reflects a strong sense of ownership and recognition of their value.

CONCLUSION AND WAY FORWARD

Water scarcity remains a pressing challenge for rural hill communities in Nepal, compounded by climate change, erratic rainfall and unplanned infrastructure development. This study has demonstrated how the EbA strategies, particularly the construction and rehabilitation of conservation ponds, can effectively address these challenges by enhancing water yield, improving soil moisture and contributing to agricultural productivity.

Community perceptions, supported by both qualitative and quantitative data, underscore the tangible benefits of these interventions. Respondents from Dolakha, Salyan and Achham districts reported significant improvements in water availability, soil moisture and greenery. Women and children, in particular, have experienced reduced workloads associated with water collection, improving their overall quality of life.

The findings highlight the importance of incorporating the EbA measures into local and national development policies as sustainable solutions to water resource management. By fostering community engagement and leveraging local knowledge, these interventions can enhance resilience against climate variability while supporting livelihoods and biodiversity.

The success of the EbA II project provides a model for replication in other vulnerable

regions, emphasising the potential for ecosystem-based strategies to mitigate climate impacts and sustainably manage water resources for the long term.

ACKNOWLEDGEMENT

The researchers extend their heartfelt gratitude to the community members of Dolakha, Salyan and Achham districts who participated in this research for generously providing their time and sharing their knowledge and experiences. Special thanks to the officials of the Climate Change Management Division, Ministry of Forests and Environment, and the project team of the EbA II Project for their guidance and support in preparing this research paper. Their comments and suggestions were invaluable for validating the research tools and enhancing the study's credibility and the reliability of its results. We also express our sincere thanks to the Global Environment Facility (GEF) and the United Nations Environment Programme (UNEP) for their financial support to the EbA II Project and this research.

REFERENCES

- CGED-Nepal.** 2024. Retrieved from <https://www.cgednepal.org/>
- FAO.** 2014. *Building a common vision for sustainable food and agriculture: Principles and approaches.* United Nations Food and Agriculture Organization, Rome, Italy.
- Garcia-Herrero, L., Lavrnic, S., Guerrieri, V., Toscano, A., Milani, M., Cirelli, G. L. and Vittuari, M.** 2022. Cost-benefit of green infrastructures for water management: A sustainability assessment of full-scale constructed wetlands in Northern and Southern Italy. *Ecological Engineering*, **185**: 1-9. <https://doi.org/10.48550/arXiv.2305.06284>.
- GFDRR.** 2021. *Vulnerability, Risk Reduction, and Adaptation to Climate Change.* Global Facility for Disaster Reduction and Recovery (GFDRR).

- Koirala, P.** 2021. *Indigenous and traditional knowledge in Nepal and its role in scaling up climate action*, viewed on 04 January 2025. <https://us.boell.org/en/2021/10/19/indigenous-and-traditional-knowledge-nepal-and-its-role-scaling-climate-action>
- Liniger, H. P., Mekdaschi Studer, R., Hauert, C. and Gurtner, M.** 2011. *Sustainable land management in practice: Guidelines and best practices for sub-Saharan Africa*. TerrAfrica, World Overview of Conservation Approaches and Technologies (WOCAT), and FAO.
- MOFE.** 2020. *Nationally Determined Contribution*. Ministry of Forests and Environment, Kathmandu, Nepal.
- MOFE.** 2023. *National Adaptation Plan for 2021–50*. Ministry of Forests and Environment, Kathmandu, Nepal.
- Secretariat of the Convention on Biological Diversity (SCBD).** 2009. *Connecting biodiversity and climate change mitigation and adaptation: Report of the second ad hoc technical expert group on biodiversity and climate change*. CBD Technical Series No. 41. Montreal: Convention on Biological Diversity.
- Shrestha, A. B., Agrawal, N. K., Alfthan, B., Bajracharya, S. R., Marechal, J. and van Oort, B. (Eds.).** 2015. *The Himalayan climate and water atlas: Impact of climate change on water resources in five of Asia's major river basins*. ICIMOD, Grid-Arendal, and CICERO.
- Thiaw, M. R., Alverson, I., Liu, J. and Han, Z.** 2013. The role of ecosystem services in climate change adaptation and disaster risk reduction. *Current Opinion in Environmental Sustainability*, 5(1): 47–52. <https://doi.org/10.1016/j.cosust.2013.02.002>.
- Triola, M. F.** 2020. *Elementary Statistics* (13th ed.). Pearson Education.

